arXiv:1401.0612v1 [nlin.SI] 3 Jan 2014

A NATURAL GEOMETRIC CONSTRUCTION UNDERLYING A CLASS OF LAX PAIRS

I.S. KRASIL′SHCHIK

Abstract. In the framework of the theory of differential coverings [2], we discuss a general geometric construction that serves the base for the so-called Lax pairs containing differentiation with respect to the spectral parameter [4]. Such kind of objects arise, for example, when studying integrability properties of equations like the Gibbons-Tsarev one [1].

Introduction

The *Gibbons-Tsarev equation*

$$z_{yy} + z_x z_{xy} - z_y z_{xx} + 1 = 0, \tag{1}$$

see, e.g., [1], arises as a reduction of the *Benney chain* and possesses many properties of integrable systems. On the other hand, it is not integrable in the exact solitonic sense (for example, the equation does not admit Hamiltonian operators). In particular, equation (1) is the compatibility condition for the system

$$\varphi_x = \frac{1}{z_y + z_x\varphi - \varphi^2}, \quad \varphi_y = -\frac{z_x - \varphi}{z_y + z_x\varphi - \varphi^2}. \tag{2}$$

Introducing a new function $\psi = \psi(\varphi)$, one can consider the system ('a Lax pair')

$$\psi_x = -\frac{1}{z_y + z_x\varphi - \varphi^2}\psi_\varphi, \quad \psi_y = \frac{z_x - \varphi}{z_y + z_x\varphi - \varphi^2}\psi_\varphi \tag{3}$$

(see, for example, [4]), whose compatibility condition is the Gibbons-Tsarev equation as well. Taking now φ for a parameter in (3) and expanding the resulting system in Laurent series, one can describe an infinite family of nonlocal conservation laws and the corresponding infinite algebra of nonlocal symmetries.

Of course, these computations can be done in a more general situation. Let $\mathscr{E}$ be a PDE imposed on an unknown function $z(x, y)$ and $\varphi_x = X(x, y, [z], \varphi)$, $\varphi_y = Y(x, y, [z], \varphi)$ be a system compatible by virtue of $\mathscr{E}$, where $[z]$ denotes a collection consisting of z itself and its derivatives up to a certain order. Then one can pass to the system $\psi_x = -X(x, y, [z], \varphi)\psi_\varphi$, $\psi_y = -Y(x, y, [z], \varphi)\psi_\varphi$ also compatible over $\mathscr{E}$. As I see it myself, this seemingly simple construction invokes two questions at least: (a) why do

2010 *Mathematics Subject Classification.* 37K10.

Key words and phrases. Geometry of PDEs, differential coverings.

I am grateful to the Mathematical Institute of the Silesian University in Opava for support and comfortable working condition.

the signs change? (b) how does the unknown function φ in the initial system transform to an independent variable in the resulting one?

Below we shall discuss a general geometric construction that (hopefully) answers these questions and explains the trick of passing from System (2) to System (3). In Section 1 we briefly recall the necessary definitions and results from the geometric theory of differential equations (see [3], for example), including the nonlocal theory, [2]. Section 2 concerns with the definition of jet spaces associated to integrable distributions. Finally, in Section 3 the main construction is described.

1. The basic notions and notation

Consider a smooth manifold M, $\dim M = n$, and a locally trivial vector fiber bundle $\pi\colon E \to M$, $\operatorname{rank}\pi = m$. Denote by $\Gamma(\pi)$ the $C^\infty(M)$-module of sections of the bundle π. Let $\pi_k\colon J^k(\pi) \to M$ be the corresponding bundle of k-jets, $k = 1, \dots, \infty$, and let $\pi_{k,l}\colon J^k(\pi) \to J^l(\pi)$, $k > l$, denote the natural projections. The bundle π_∞ possesses a natural flat connection $\mathscr{C}$, which is called the *Cartan connection* and is defined by the condition

$$j_\infty(f)^*(\mathscr{C}_X(\varphi)) = X(j_\infty(f)^*(\varphi)),$$

where X is a vector field on M, φ is a smooth function on $J^\infty(\pi)$, f is a section of π, and $j_\infty(f) \in \Gamma(\pi_\infty)$ denotes the infinite jet of this section. The corresponding horizontal distribution $z \mapsto \mathscr{C}_z$, $z \in J^\infty(\pi)$, on $J^\infty(\pi)$ is integrable and is called the *Cartan distribution*.

An (*infinitely prolonged*) *differential equation* will be understood as a submanifold $\mathscr{E} \subset J^\infty(\pi)$ such that all the fields of the form $\mathscr{C}_X$ are tangent to this submanifold. Consequently, the Cartan connection can be restricted to the projection $\pi_\infty|_{\mathscr{E}}$, while the Cartan distribution is restricted to $\mathscr{E}$. We shall keep the same notation for these restrictions.

Let $\tilde{\mathscr{E}} \subset J^\infty(\tilde\pi)$, $\tilde\pi\colon \tilde E \to M$, be another equation. A smooth vector bundle $\tau\colon \tilde{\mathscr{E}} \to \mathscr{E}$ is called a (*differential*) covering if one has $\tau_* \circ \tilde{\mathscr{C}}_X = \mathscr{C}_X$ for any vector field $X \in D(M)$, where $D(M)$ denotes the module of vector fields on M and 'star' stands for the differential of a smooth map.

Coordinates. Let $\mathscr{U} \subset M$ be a local chart in M with coordinates $x^1, \dots, x^n$ and $\pi^{-1}(\mathscr{U}) = \mathscr{U} \times \mathbb{R}^m$ be a trivialization of the bundle π with some coordinates $u^1, \dots, u^m$ in the fibers. Then the *adapted coordinates* u^j_σ arise in $J^\infty(\pi)$. The Cartan connection in these coordinates is defined by the *total derivative* operators

$$D_{x^i} = \frac{\partial}{\partial x^i} + \sum_{j,\sigma} u^j_{\sigma i} \frac{\partial}{\partial u^j_\sigma},$$

where σ is the multi-index corresponding to the variables x^i. A covering structure in the bundle $\tau\colon \tilde{\mathscr{E}} \to \mathscr{E}$ with coordinates $w^1, \dots, w^r$ along the fibers (the functions w^γ are called *nonlocal variables*) is given by the extensions of the total derivatives

$$\tilde D_{x^i} = D_{x^i} + X_i, \tag{4}$$

where X_i are τ-vertical vector fields (i.e., the fields of the form $\sum_\gamma X_i^\gamma \partial/\partial w^\gamma$) that satisfy the relations

$$D_{x^i}(X_j) - D_{x^j}(X_i) + [X_i, X_j] = 0$$

for all $i < j$.

A covering τ is called *trivial* if $\mathscr{C}_X(C^\infty(\tilde{\mathscr{E}})) \subset C^\infty(\mathscr{E})$ for all vector fields $X \in D(M)$. A *morphism* of two coverings $\tau_i \colon \tilde{\mathscr{E}}_\alpha \to \mathscr{E}$, $\alpha = 1, 2$, of ranks r_1 and r_2, respectively, is a smooth map $\varphi \colon \tilde{\mathscr{E}}_1 \to \tilde{\mathscr{E}}_2$ such that

(1) $\tau_1 = \tau_2 \circ \varphi$ and
(2) $\varphi_*(\tilde{\mathscr{C}}^1_z) \subset \tilde{\mathscr{C}}^2_{\varphi(z)}$, $z \in \tilde{\mathscr{E}}$.

Coordinates. A covering τ is trivial if and only if the vertical fields in Equalities (4) vanish.

If w^j_α are the nonlocal variables in the covering τ_α, $\alpha = 1, 2$, while the covering structures are given by the vertical fields

$$\sum_\gamma X_i^{\alpha,\gamma} \frac{\partial}{\partial w^\gamma_\alpha}$$

then a morphism φ of τ_1 to τ_2 is determined by a system of functions $w_1^\gamma = \varphi^\gamma(w_1^1, \dots, w_1^{r_1})$, $\gamma = 1, \dots, r_2$, such that $\tilde{D}^1_{x^i}(\varphi^\gamma) = \varphi^*(X_i^{2,\gamma})$. In particular, φ is a morphism to the trivial covering if

$$\tilde{D}_{x^i}(\varphi^\gamma) = 0. \tag{5}$$

Such morphisms will be called *trivializing* ones.

2. Jets associated to distributions

Consider a smooth manifold N, $\dim N \leq \infty$. Let $\mathscr{D}$ be an integrable distribution on this manifold, $\operatorname{rank} \mathscr{D} = s < \infty$, i.e., a locally projective finite-rank submodule $\mathscr{D} \subset D(N)$ such that $[\mathscr{D}, \mathscr{D}] \subset \mathscr{D}$. Let also $\xi \colon F \to N$ be a locally trivial vector bundle over N. We say that two sections f and $f' \in \Gamma(\xi)$ are *k-equivalent* over $\mathscr{D}$ at a point $z \in N$ if

$$\big(X_1 \dots X_l(f)\big)\big|_z = \big(X_1 \dots X_l(f')\big)\big|_z, \qquad l \leq k,$$

for any collection of vector fields $X_1, \dots, X_l \in \mathscr{D}$. Let $[f]^k_z$ denote the equivalence class of f. The set

$$J^k_{\mathscr{D}}(\xi) = \{\, [f]^k_z \mid z \in N, f \in \Gamma(\xi) \,\}, \quad k = 0, 1, \dots, \infty,$$

is endowed with a natural smooth manifold structure (cf. [3]) and is called the *the manifold of k-jets associated to the distribution $\mathscr{D}$*. One has the natural projections $\xi_k \colon J^k_{\mathscr{D}}(\xi) \to M$ and $\xi_{k,l} \colon J^k_{\mathscr{D}}(\xi) \to J^l_{\mathscr{D}}(\xi)$, $k > l$, and all these maps are locally trivial fiber bundles while the functions $j_k(f) \colon M \to J^k_{\mathscr{D}}(\xi)$, $z \mapsto [f]^k_z$, (*jets of sections*) are smooth sections of the bundles ξ_k.

Example. If $\mathscr{D} = D(N)$ we obtain the classical definition of jet spaces.

Example. If $N = \mathscr{E}$ is an equation and $\mathscr{D} = \mathscr{C}$ coincides with the Cartan distribution we arrive to the definition of *horizontal jets*, see [3].

The bundle of infinite jets ξ_∞ admits a flat $\mathscr{D}$-*connection* denoted by

$$\nabla^{\mathscr{D}}\colon \mathscr{D} \to D(J^\infty_{\mathscr{D}}(\xi))$$

and which is also called the *Cartan connection*: for any point $\theta = [f]^\infty_z$, a field $X \in \mathscr{D}$ and a function $\varphi \in C^\infty(J^\infty_{\mathscr{D}}(\xi))$, we set

$$\nabla^{\mathscr{D}}_X(\varphi)\Big|_\theta = X(j_\infty(f)^*\varphi)|_z\,.$$

The corresponding horizontal distribution on $J^\infty_{\mathscr{D}}(\xi)$ will be denoted by $\Delta^{\mathscr{D}}$; it is integrable, i.e., $[\Delta^{\mathscr{D}}, \Delta^{\mathscr{D}}] \subset \Delta^{\mathscr{D}}$.

A submanifold $\mathscr{E}_k \subset J^k_{\mathscr{D}}(\xi)$ is called a $\mathscr{D}$-*equation* of order k. As in the case of 'usual' equations, one can define the $\mathscr{D}$-*prolongations* $\mathscr{E}_k^{(l)} \subset J^{k+l}_{\mathscr{D}}(\xi)$ of finite and infinite orders. The latter, $\mathscr{E} \subset J^\infty_{\mathscr{D}}(\xi)$, will be shortly called just a $\mathscr{D}$-*equation*. The map $\xi_\infty\colon \mathscr{E} \to M$ inherits the Cartan connection, while the submanifols $\mathscr{E}$ carries the corresponding horizontal distribution.

Let S be a ξ_∞-vertical vector field on $\mathscr{E}$; it is called a (*higher infinitesimal*) $\mathscr{D}$-*symmetry* of the equation $\mathscr{E}$ if $[S, \Delta^{\mathscr{D}}] \subset \Delta^{\mathscr{D}}$.

Coordinates. Let $y^1, \dots, y^k, \dots$ be local coordinates in N, $v^1, \dots, v^r$ be coordinates if fibers of a trivialization of the bundle ξ. Finally, let $Y_1, \dots, Y_s$ be a local basis of the distribution $\mathscr{D}$,

$$[Y_i, Y_j] = \sum_k c^k_{ij} Y_k, \qquad c^k_{ij} \in C^\infty(J^\infty_{\mathscr{D}}(\xi)).$$

Let $\sigma = i_1 \dots i_{|\sigma|}$ be a multi-index of arbitrary but finite length $|\sigma|$, $1 \leq i_\alpha \leq s$. Let us define the coordinate functions v^j_σ on $J^\infty_{\mathscr{D}}(\xi)$ by setting

$$v^j_\sigma(\theta) = Y_{i_1} \dots Y_{i_{|\sigma|}}(f^j)\Big|_z\,, \qquad \theta = [f]^\infty_z, \quad f = (f^1, \dots, f^l).$$

Then the lifts $\nabla^{\mathscr{D}}_Y$ of vector fields $Y \in \mathscr{D}$ to $J^\infty_{\mathscr{D}}(\xi)$, i.e., the connection $\nabla^{\mathscr{D}}$, are defined by the equalities

$$\nabla^{\mathscr{D}}_{Y_\alpha}(v^j_\sigma) = v^j_{\alpha\sigma}.$$

One has the following relations

$$v^j_{\alpha\sigma} = v^j_{i_1\alpha\bar\sigma} + \sum_k c^k_{\alpha i_1} v^j_{k\bar\sigma}, \tag{6}$$

where $\bar\sigma = i_2 \dots i_{|\sigma|}$. The functions

$$v^j_K = v_{\underbrace{1\dots1}_{k_1\text{times}}\dots\underbrace{s\dots s}_{k_s\text{times}}}.$$

may be taken for independent coordinates.

Obviously, $\mathscr{D}$-*coverings* of $\mathscr{D}$-equations are also defined in a straightforward way.

3. Jets over differential coverings

Let us now consider the constructions of Section 2 when $N = \tilde{\mathscr{E}}$, where $\tilde{\mathscr{E}}$ is the covering equation in some covering $\tau\colon \tilde{\mathscr{E}} \to \mathscr{E}$. Then the manifold $\tilde{\mathscr{E}}$ carries three integrable distributions:

- the *horizontal distribution* denoted by $\mathscr{H}$ which coincides with the Cartan distribution $\tilde{\mathscr{C}}$ on $\tilde{\mathscr{E}}$;
- the *vertical distribution* $\mathscr{V}$ generated by τ-vertical vector fields;
- the *total distribution* $\mathscr{T}$ which is the sum of the previous two.

Respectively, one can define the jet spaces associated to these distributions and the diagram



is commutative (the projections $\xi^{\mathscr{T}\mathscr{H}}$ and $\xi^{\mathscr{T}\mathscr{V}}$ are defined in an obvious way).

Note that the distribution $\Delta^{\mathscr{T}}$, just like the original total distribution $\mathscr{T}$, is the sum of the distributions $\Delta^{\mathscr{H}}$ and $\Delta^{\mathscr{V}}$, where $\Delta^{\mathscr{H}}$ and $\Delta^{\mathscr{V}}$ are obtained by lifting $\mathscr{H}$ and $\mathscr{V}$ using the Cartan connection associated to the distribution $\mathscr{T}$. Thus, the jet manifold $J^\infty_{\mathscr{T}}(\xi)$ possesses *three different geometries*. Note also that the projection $\xi^{\mathscr{T}\mathscr{V}}$ is a covering with respect to the $\mathscr{V}$-geometry while the $\xi^{\mathscr{T}\mathscr{H}}$ is a covering in the $\mathscr{H}$-geometry of the space $J^\infty_{\mathscr{T}}(\xi)$.

Note also that for any $\mathscr{H}$-equation $\mathscr{W} \subset J^\infty_{\mathscr{H}}(\xi)$ one can consider its preimage $(\xi^{\mathscr{T}\mathscr{H}})^{-1}(\mathscr{W})$ in $J^\infty_{\mathscr{T}}(\xi)$ and the infinite $\mathscr{T}$-prolongation of the latter. The resulting $\mathscr{T}$-equation is denoted by $\tilde{\mathscr{W}}$ also possesses three different geometries, while the map $\xi^{\mathscr{T}\mathscr{H}}\big|_{\tilde{\mathscr{W}}}$ is a covering in the geometry associated to the horizontal distribution. Exactly the same situation arises when one considers $\mathscr{V}$-equations.

Coordinates. Let the covering structure in the bundle $\tau\colon \tilde{\mathscr{E}} \to \mathscr{E}$ be defined by vector fields (4) and choose

$$\frac{\partial}{\partial w^1}, \dots, \frac{\partial}{\partial w^r}$$

for a basis in the space of τ-vertical fields. Then

$$[\tilde{D}_{x^i}, \tilde{D}_{x^j}] = 0, \quad [\frac{\partial}{\partial w^\alpha}, \frac{\partial}{\partial w^\beta}] = 0, \quad [\frac{\partial}{\partial w^\alpha}, D_{x^i}] = \sum_\beta \frac{\partial X^\beta_i}{\partial w^\alpha} \frac{\partial}{\partial w^\beta}. \qquad (7)$$

Denote by σ the multi-index that corresponds to the variables x^i. Let also the multi-index ρ correspond to the variables w^α. Then we obtain the following coordinate functions in the jet spaces:

- v^j_σ in $J^\infty_{\mathscr{H}}(\xi)$;

- v^j_ρ in $J^\infty_{\mathscr{V}}(\xi)$;
- $v^j_{\sigma\rho}$ in $J^\infty_{\mathscr{T}}(\xi)$.

In these coordinates, one has

$$\Delta^{\mathscr{T}}_{x^i} = \tilde{D}_{x^i} + \sum v^j_{i\sigma\rho}\frac{\partial}{\partial v^j_{\sigma\rho}}, \quad \Delta^{\mathscr{T}}_{w^\alpha} = \frac{\partial}{\partial w^\alpha} + \sum v^j_{\alpha\sigma\rho}\frac{\partial}{\partial w^\alpha},$$

while the functions $v^j_{i\sigma\rho}$, $v^j_{\alpha\sigma\rho}$ are computed accordingly to Equations (6) and (7).

Existence of different geometries on the manifold $J^\infty_{\mathscr{T}}(\xi)$ leads to different notions of a symmetry. Below, we shall need the following one: *a $\xi^{\mathscr{T}}_\infty$-vertical field S is called a gauge $\mathscr{V}$-symmetry if $[S, \Delta^{\mathscr{V}}] \subset \Delta^{\mathscr{V}}$*. Similar to the classical case, one has the following description of these symmetries:

Proposition 1. *Gauge $\mathscr{V}$-symmetries are in one-to-one correspondence with sections of the pull-back $(\xi^{\mathscr{T}}_\infty)^*(\xi^{\mathscr{H}}_\infty)$. In local coordinates, to any such a section h of the form $v^j_\sigma = h^j_\sigma(\theta)$, $\theta \in J^\infty_{\mathscr{T}}(\xi)$, there corresponds the symmetry*

$$\mathbf{E}_h = \sum_{\rho,\gamma,j} \Delta^{\mathscr{T}}_\rho(h^j_\sigma)\frac{\partial}{\partial v^j_{\rho\sigma}},$$

where $\Delta^{\mathscr{T}}_\rho$ is the composition of the total derivatives $\Delta^{\mathscr{T}}_{w^\alpha}$ corresponding to the multi-index ρ.

If $\mathscr{W} \subset J^\infty_{\mathscr{T}}(\xi)$ is a $\mathscr{T}$-equation then the Lie algebra of its gauge $\mathscr{V}$-symmetries will be denoted by $\mathfrak{gs}(\mathscr{W})$.

4. Jets of automorphisms

Consider now a particular case of the above described constructions: let $\xi = \tau^*(\tau)$. Sections of this bundle are naturally identified with morphisms of the bundle τ, i.e., with smooth maps $\varphi\colon \tilde{\mathscr{E}} \to \tilde{\mathscr{E}}$, such that $\tau \circ \varphi = \tau$.

Consider the submanifold

$$\overline{J^\infty_{\mathscr{H}}(\xi)} = \{\, \theta \in J^\infty_{\mathscr{H}}(\xi) \mid \theta = [\varphi]^\infty_z,\ \varphi(z) = z,\ z \in \tilde{\mathscr{E}}\,\} \subset J^\infty_{\mathscr{H}}(\xi)$$

in $J^\infty_{\mathscr{H}}(\xi)$ and the $\mathscr{H}$-equation

$$\mathscr{W}(\tau) = \{\, \theta \in \overline{J^\infty_{\mathscr{H}}(\xi)} \mid \theta = [\varphi]^\infty_z,\ \varphi \text{ is a trivializing morphism}\,\}.$$

Let $\tilde{\mathscr{W}}(\tau) \subset J^\infty_{\mathscr{T}}(\xi)$ be the $\mathscr{T}$-prolongation of this equation endowed with the $\mathscr{H}$-geometry, i.e., equipped with the horizontal distribution, as it was described above. Then we obtain the infinite-dimensional covering

$$\tilde{\tau}\colon \tilde{\mathscr{W}}(\tau) \to \mathscr{E}, \tag{8}$$

which is associated with the initial covering τ in a canonical way.

Coordinates. Let φ^α_σ be local coordinates in $\overline{J^\infty_{\mathscr{H}}(\xi)}$, where σ is the multi-index corresponding to the variables $x^1, \dots, x^n$. Then the equation $\mathscr{W}(\tau)$ is determined by the relations

$$\varphi^\gamma_{x^i} \equiv \tilde{D}_{x^i}(\varphi^\gamma) = 0, \qquad i = 1, \dots, \dim M, \quad \gamma = 1, \dots, \operatorname{rank}\tau,$$

or

$$\frac{\partial\varphi^\gamma}{\partial x^i}+\sum_\alpha X_i^\alpha\frac{\partial\varphi^\gamma}{\partial w^\alpha}=0. \tag{9}$$

The nonlocal variables in the covering $\tilde\tau$ are the functions $\varphi^\gamma_\rho=\Delta^{\mathscr{T}}_\rho(\varphi^\gamma)$, where ρ is the multi-index that corresponds to the variables $w^1,\dots,w^r$. Then the equation $\tilde{\mathscr{W}}(\tau)$ is given by the infinite system

$$\frac{\partial\varphi^\gamma_\rho}{\partial x^i}+\Delta^{\mathscr{T}}_\rho\Big(\sum_\alpha X_i^\alpha\varphi^\gamma_\alpha\Big)=0.$$

Thus, the covering structure in the bundle $\tilde\tau$ is described by the vector fields

$$\Delta^{\mathscr{T}}_{x^i}=D_{x^i}+\sum_{\rho,\gamma}\Delta^{\mathscr{T}}_\rho\Big(\sum_\alpha X_i^\alpha\varphi^\gamma_\alpha\Big)\frac{\partial}{\partial\varphi^\gamma_\rho},$$

or

$$\Delta^{\mathscr{T}}_{x^i}=D_{x^i}+\mathbf{E}_{h_i}, \tag{10}$$

where

$$h_i=\left(\sum_\alpha X_i^\alpha\varphi^1_\alpha,\dots,\sum_\alpha X_i^\alpha\varphi^r_\alpha\right),\quad i=1,\dots,n.$$

Summarizing the above discussion, we arrive to the following

Proposition 2. *The covering $\tilde\tau\colon\tilde{\mathscr{W}}(\tau)\to\mathscr{E}$ defined by equalities* (10) *is a zero-curvature representation with values in the Lie algebra of gauge $\mathscr{V}$-symmetries $\mathfrak{gs}(\mathscr{W})$.*

Obviously, the construction introduced here is a geometrical generalization of the 'Lax pairs' mentioned in Introduction.

Acknowledgment

I am grateful to Dr. Michal Marvan for discussions and fruitful cooperation.

References

[1] J. Gibbons, S.P. Tsarev. *Reductions of the Benney Equations*. Phys. Letters A., **211** (1996) no. 1, pp. 19–24.

[2] I.S. Krasil'shchik, A.M. Vinogradov. *Nonlocal trends in the geometry of differential equations: Symmetries, conservation laws, and Bäcklund transformations*. Acta Applicandae Mathematica, **15** (1989) no. 1-2, pp. 161–209.

[3] I.S. Krasil′shchik, V.V. Lychagin, and A. M. Vinogradov, *Geometry of jet spaces and nonlinear partial differential equations*, Gordon and Breach, 1986.

[4] A.V. Odesskii, V.V. Sokolov. *Non-homogeneous systems of hydrodynamic type possessing Lax representations*. Commun. Math. Phys., **324** (2013) no. 1, pp. 47–62.

Slezská univerzita v Opavě, Matematický ústav v Opavě, Na Rybníčku 626/1, 746 01 Opava, Czech Republic & Independent University of Moscow, 119002, B. Vlasyevskiy Per. 11, Moscow, Russia

E-mail address: josephkra@gmail.com